

---

**Human germ cell formation in xenotransplants of induced pluripotent stem cells carrying X chromosome aneuploidies.**

**Journal:** Sci Rep

**Publication Year:** 2014

**Authors:** Antonia A Dominguez, H Rosaria Chiang, Meena Sukhwani, Kyle E Orwig, Renee A Reijo Pera

**PubMed link:** 25242416

**Funding Grants:** Stanford CIRM Training Program

**Public Summary:**

**Scientific Abstract:**

Turner syndrome is caused by complete or partial loss of the second sex chromosome and is characterized by spontaneous fetal loss in >90% of conceptions. Survivors possess an array of somatic and germline clinical characteristics. Induced pluripotent stem cells (iPSCs) offer an opportunity for insight into genetic requirements of the X chromosome linked to Turner syndrome. We derived iPSCs from Turner syndrome and control individuals and examined germ cell development as a function of X chromosome composition. We demonstrate that two X chromosomes are not necessary for reprogramming or maintenance of pluripotency and that there are minimal differences in gene expression, at the single cell level, linked to X chromosome aneuploidies. Formation of germ cells, as assessed in vivo through a murine xenotransplantation model, indicated that undifferentiated iPSCs, independent of X chromosome composition, are capable of forming germ-cell-like cells (GCLCs) in vivo. In combination with clinical data regarding infertility in women with X chromosome aneuploidies, results suggest that two intact X chromosomes are not required for human germ cell formation, qualitatively or quantitatively, but rather are likely to be required for maintenance of human germ cells to adulthood.

---

**Source URL:** <https://www.cirm.ca.gov/about-cirm/publications/human-germ-cell-formation-xenotransplants-induced-pluripotent-stem-cells>